

STIC-ILL

*Mic
RC583.C45
Adonis*

From: Huynh, Phuong N.
Sent: Tuesday, January 30, 2001 6:00 PM
To: STIC-ILL
Subject: RE: 09/270,910 Rush

Hi!

Please deliver the following references:

1. Prot Exp Purification 8: 365-373 (1996).
2. Clin. Exp. Allergy 22: 391-9 (1992)
3. Nature structural biology 3: 1040-45 (1996).

Thanks,
Neon
Art unit 1644
Mail CM1 9E12
Tel 308-4844

Specificity mapping of patients IgE response towards the tree pollen major allergens *Aln g I*, *Bet v I* and *Cor a I*

H. IPSEN*, J.-Å. WIHL†, B. NÜCHEL PETERSEN‡ and H. LØWENSTEIN*

*ALK Research, Hørsholm, Denmark, †Department of Otorhinolaryngology, University of Lund, Malmö General Hospital, Sweden and ‡Department of Lung Diseases, Copenhagen County Hospital, Gentofte, Denmark

Summary

The specificity pattern of IgE from non-treated tree pollen allergic patients ($n = 38$) were evaluated by solid phase absorption of serum samples followed by CRIE on alder, birch and hazel CIE precipitation profiles. The majority of the serum samples seemed to contain IgE antibodies with the following characteristics: specific towards *Bet v I* alone and common between *Aln g I*, *Bet v I* and/or *Cor a I*. The IgE specificity profiles observed for 95% of the sera tested are compatible with birch pollen allergens being the only sensitizing allergens, indicating that the patients react to allergens from other tree pollens of the Fagales order due to IgE cross-reaction with the major allergens of birch and alder and/or hazel pollens.

Clinical and Experimental Allergy, Vol. 22, pp. 391–399. Submitted 16 May 1991; revised 21 October 1991; accepted 25 October 1991.

Introduction

Patients allergic towards a particular allergen source often react towards the taxonomically related species of the same genus, e.g. tree (birch) pollen sensitive patients react towards pollen from many species of the Fagales order [1–6]; grass pollen sensitive patients react towards many of the Poales species [1,7,8] and house dust mite sensitive patients often react towards all three dermatophagoides species [9–12]. Indeed, the major allergens from taxonomically related species exhibit a high degree of similarity both with respect to physico-chemical parameters and immunochemical reactivity towards both poly- and monoclonal antibodies [7,10,12–35]. An increasing number of major allergens have been sequenced completely or in part either by conventional methods or by recombinant DNA techniques [10,23,25,29,34–45], and the amino acid sequences of allergens from taxonomically related species often exhibit 60–70% sequence identity [10,29,33–41,43] and still higher degrees of sequence homology. This indicates that allergens from related species presumably exhibit epitope patterns which are likely to be more or less preserved within the genus. Therefore, it is probable that the patients concordant sensitivity towards species related allergens originate

from congruent or partial identical epitopes. The present paper describes the classification of the specificity of IgE from birch pollen sensitive patients with respect to the cross reaction towards pollen major allergens from the related species alder and hazel.

Materials and methods

Pollen extracts

Aqueous extracts of alder (*Alnus glutinosa*), birch (*Betula verrucosa*) and hazel (*Corylus avellana*) pollens were prepared and handled as described recently [19,29].

Rabbit antibodies

Rabbit anti-alder (a-Ag), anti-birch (a-Bv) and anti-hazel (a-Ca) pollen extract antibodies were raised as described [19,29].

Patient sera

Serum samples ($n = 38$) from non-treated patients with seasonal allergy towards pollens from trees of the Fagales order were selected as described in detail elsewhere [2–4].

Immunoelectrophoretic methods

Crossed immunoelectrophoresis [20,29,46] was performed in 1% (w/v) agarose (type HSA, Litex, Denmark)

Correspondence: Dr H. Ipsen, ALK Research, Bøge Allé 10–12, DK-2970, Hørsholm, Denmark.

(0.073 M Tris, 0.024 M barbitale, 0.0006 M calcium lactate pH 8.6) moulded on 5 by 7 cm glass plates. Extracts of alder (Ag, 10 µg), birch (Bv, 15 µg) and hazel (Ca, 10 µg) were applied into wells punched in the agarose at the side measuring 7 cm. First dimension electrophoresis: 15 min for Ag and Bv, 20 min for Ca at 10 V/cm, 15°C; second dimension electrophoresis: 16 hr at 2 V/cm 15°C into separately moulded agarose gels containing 12.5 µl/cm² a-Ag, 16 µl/cm² a-Bv and 11.1 µl/cm² a-Ca rabbit antibodies, respectively. Cold dried crossed immuno-

plates were incubated with either patient serum (diluted 1:40 (v/v) in 0.027 M Na₂HPO₄, 0.019 M NaH₂PO₄, I=0.1, pH 7.0) (8 ml) or solid phase absorbed sera (8 ml) (see below) for 16 hr at room temperature, washed three times 10 min each with the dilution buffer and incubated with ¹²⁵I-anti-IgE [18,47] (approximately 125 000 c.p.m.) in 0.027 M Na₂HPO₄, 0.019 M NaH₂PO₄, 0.3% (w/v) bovine serum albumin, 0.154 M NaCl, 0.015 M NaN₃, 0.027 M EDTA pH 7.5 for 16 hr at room temperature. After three 10 min washes (0.154 M NaCl, 1% (v/v) Tween 20), the

Table 1. Listing of the specificities which can account for a radiostaining of a given major allergen. The columns list the specificity combinations allowed after a given solid phase absorption. A, B, C1 and C2 signifies radiostaining of *Aln g I*, *Bet v I*, *Cor a I* and *Cor a 'II'* respectively. Combinations of lettering signifies concordant reactivity

	Ø-Sepharose	Ag-Sepharose	Bv-Sepharose	Ca-Sepharose
<i>Aln g I</i>	ABC1C2 ABC1 ABC2 AC1C2 AB AC1 AC2 A		AC1C2 AC1 AC2 A	AB A
<i>Bet v I</i>	ABC1C2 ABC1 ABC2 BC1C2 AB BC1 BC2 B	BC1C2 BC1 BC2 B		AB B
<i>Cor a I</i>	ABC1C2 ABC1 AC1C2 BC1C2 AC1 BC1 C1C2 C1	BC1C2 BC1 C1C2 C1	AC1C2 AC1 C1C2 C1	
<i>Cor a 'II'</i>	ABC1C2 ABC2 AC1C2 BC1C2 AC2 BC2 C1C2 C2	BC1C2 BC2 C1C2 C2	AC1C2 AC2 C1C2 C2	

crossed immunoplates were dried and exposed on Kodak XL-1 X-ray film for 3 and 10 days at -20°C using an intensifying screen. SRRID was performed as described elsewhere [48].

Solid phase absorption of patient sera

The solid phase absorption experiments were performed in a manner similar to those described by Lind *et al.* [9].

The extracts Ag, Bv and Ca were each covalently coupled to CNBr activated Sepharose CL 4B (Pharmacia, Sweden) according to the manufactures' recommendations (20 mg extract per g of CNBr activated Sepharose CL 4B).

Solid phase absorption was performed with 38 sera. One millilitre of patient serum was applied on a column containing 2 ml Protein A Sepharose CL 4B (Pharmacia, Sweden), equilibrated with 0.027 M Na_2HPO_4 , 0.019 M NaH_2PO_4 , $\text{I}=0.1$, pH 7.0 and the column was eluted with 7 ml of the same buffer. The IgG depleted patient serum (column eluate, approximately 8 ml) was aliquoted into four 2 ml samples and applied on columns containing 1 ml of either Sepharose CL 4B, Ag-, Bv- or Ca-Sepharose CL 4B equilibrated with 0.027 M Na_2HPO_4 , 0.019 M NaH_2PO_4 , $\text{I}=0.1$, pH 7.0. The serum was eluted from each of the extract-Sepharose columns with 6 ml 0.027 M Na_2HPO_4 , 0.019 M NaH_2PO_4 , $\text{I}=0.1$, pH 7.0 and the eluates (8 ml) were each incubated with crossed immunoplates as described above. A set of solid phases were used for one serum sample only.

Quantitation of the IgE binding

The radiostaining intensity on the autoradiographs of the images of the major allergen (*Aln g I*, *Bet v I*, *Cor a I* and 'II') precipitates was determined by visual comparison with the intensity of the images of the radiostaining on autoradiographs of several (nine) dilutions of ^{125}I -anti-IgE precipitated in SRRID rings [48]. Each radiostained major allergen was assigned a score number, one through nine, corresponding to the SRRID ring exhibiting a radiostaining with approximately equal intensity. The score number represents twofold dilutions of ^{125}I -anti-IgE, where nine is undiluted and one represents a 512 fold dilution.

Theoretical considerations of cross-reactivity

The apparent specificity of the IgE antibodies towards the major allergens may be separated into two different general reactivity patterns: apparent distinct and apparent common specificity. Distinct specificity represents IgE reactivity towards a particular major allergen only and common specificity represents IgE reactivity towards more than one major allergen. Given four major allergens

Aln g I (A), *Bet v I* (B), *Cor a I* (C1) and *Cor a 'II'* (C2) a total of 15 different specificity patterns can be anticipated: A; B; C1; C2; ABC1C2; ABC1; ABC2; AC1C2; BC1C2; AB; AC1; AC2; BC1; BC2 and C1C2 where a single letter represents a distinct specificity and a letter combination represents a common specificity towards the allergens represented by the letters. Any IgE reactivity representing a subset of the specificity patterns are possible. Table 1 lists the possible subsets which can account for an observed radiostaining of the individual major allergens by the IgE present in the serum sample after a given solid phase absorption. Each serum was assigned a specificity pattern by comparing Table 1 with the CRIE radiostaining profile, of the individual major allergens, obtained with the solid-phase absorbed samples.




Fig. 1. Crossed immunoelectrophoresis (CIE) precipitation pattern (a) and crossed radioimmuno-electrophoresis (CRIE) autoradiograph (10 days) (b) of alder (Ag), birch (Bv) and hazel (Ca) pollen extracts precipitated with rabbit antibodies raised towards the individual extracts. The major allergens are indicated by arrows on both the CIE and CRIE autoradiograph.

Results

The identity and position of the major allergens *Aln g I*, *Bet v I*, *Cor a I* and *Cor a II* on the CIE plates and CRIE autoradiographs are indicated on Fig. 1. The precipitates of the major allergens are all anodic and although the *Cor a II* precipitate is stained weakly on the CIE immunoplates all the major allergens were clearly visible on the CRIE autoradiographs.

The solid phase absorption procedure for the patient sera is illustrated in Fig. 2, a new set of solid phases (five) were used for each serum sample processed. Examples of the CRIE autoradiographs obtained with solid phase absorbed serum samples are shown in Fig. 3. The CRIE radiostaining intensities and reactivity patterns obtained with untreated serum and serum passed through Protein A- and plain (Ø)-Sephacrose CL 4B did not differ

significantly (Fig. 3) indicating that the column processing of the sera neither alters the specificity nor the reactivity of the IgE antibodies. Table 2 shows the CRIE score matrixes determined for all ($n=38$) the absorbed serum samples.

The score matrixes can be divided into nine different groups, each representing a unique specificity subset, which occur at widely different frequencies from 16 to 1 out of 38. The specificity pattern of the individual patients IgE was evaluated by comparing the radiostaining of the individual CRIE autoradiographs with the specificity matrix (Table 1), e.g. Fig. 3 and Table 2 serum sample number 8, Table 1. There is a clear radiostaining of *Aln g I*, *Bet v I*, *Cor a I* and *Cor a II* (Fig. 3, Ø-Sephacrose, Table 2, column Ø), initially allowing all the specificities tabulated in column Ø, Table 1. Bv-Sephacrose abolishes all radiostaining excluding the specificity patterns listed in




Fig. 2. Schematic representation of the solid phase absorption experiments.

column B, Table 1. Ag-Sepharose allows the radiostaining of *Bet v* I only, excluding the specificity patterns listed in column A, Table 1 for *Cor a* I and *Cor a* 'II' indicating distinct B specificity. Ca-Sepharose allows radiostaining of *Aln g* I and *Bet v* I verifying the existence of distinct B and further indicate common AB; distinct A is excluded by the missing radiostaining in column B, row *Aln g* I of Table 1. All the solid phases except Ø-Sepharose removes

the radiostaining of *Cor a* I and *Cor a* 'II' indicating IgE with concordant reaction towards all the major allergens. The intensity of the radiostaining (Table 2, serum sample number 8) column Ø > column C > column A further substantiates the presence of common AB but does not exclude common specificity patterns between three or four allergens, leaving serum sample number 8 with the following pattern: B, AB, ABC1, ABC2, ABC1C2.




Fig. 3. Examples of CRIE autoradiographs of patient serum samples (numbers 8, 14, 15 and 30) before and after solid phase absorption. The autoradiographs are marked to indicate the 'treatment' of the serum sample. (a) Untreated serum. (b) Ø-Sepharose. (c) Ag-Sepharose. (d) Bv-Sepharose. (e) Ca-Sepharose.

Table 2. CRIE radiostaining intensity of *Aln* g I, *Bet* v I, *Cor* a I and *Cor* a 'II' determined with solid phase absorbed serum samples ($n = 38$). Ø, plain Sepharose. A, Ag-Sepharose. B, Bv-Sepharose. C, Ca-Sepharose.

	#01	Ø	A	B	C	#02	Ø	A	B	C	#03	Ø	A	B	C	#04	Ø	A	B	C
Alng l		6	0	0	0		5	0	0	5		6	0	0	4		6	0	4	5
Bet v l		7	5	0	5		6	0	0	5		7	5	0	6		8	6	0	7
Cor a l		4	0	0	0		0	0	0	0		6	0	0	0		5	0	0	0
Cor a 'll'		5	0	0	0		0	0	0	0		6	0	0	0		6	0	0	0
	#05	Ø	A	B	C	#06	Ø	A	B	C	#07	Ø	A	B	C	#08	Ø	A	B	C
Alng l		6	0	0	5		6	0	0	4		6	0	0	5		6	0	0	6
Bet v l		7	5	0	6		7	6	0	6		7	5	0	6		7	6	0	7
Cor a l		5	0	0	0		6	0	0	0		6	0	0	0		5	0	0	0
Cor a 'll'		6	0	0	0		6	0	0	0		6	0	0	0		5	0	0	0
	#09	Ø	A	B	C	#10	Ø	A	B	C	#11	Ø	A	B	C	#12	Ø	A	B	C
Alng l		6	0	0	6		5	0	0	5		6	0	0	5		5	0	0	4
Bet v l		6	0	0	6		6	6	0	6		7	6	0	6		5	5	0	5
Cor a l		0	0	0	0		5	5	0	0		6	0	0	0		0	0	0	0
Cor a 'll'		5	0	0	0		5	5	0	0		6	0	0	0		0	0	0	0
	#13	Ø	A	B	C	#14	Ø	A	B	C	#15	Ø	A	B	C	#16	Ø	A	B	C
Alng l		5	0	0	5		4	0	0	4		5	0	0	5		7	0	4	6
Bet v l		6	5	0	6		6	0	0	5		7	6	0	7		7	6	0	7
Cor a l		4	0	0	0		0	0	0	0		0	0	0	0		7	0	0	0
Cor a 'll'		4	0	0	0		0	0	0	0		0	0	0	0		7	0	0	0
	#17	Ø	A	B	C	#18	Ø	A	B	C	#19	Ø	A	B	C	#20	Ø	A	B	C
Alng l		6	0	0	4		6	0	0	6		5	0	0	4		6	0	0	4
Bet v l		6	0	0	4		6	4	0	6		6	0	0	5		7	6	0	6
Cor a l		3	0	0	0		4	0	0	0		0	0	0	0		5	0	0	0
Cor a 'll'		4	0	0	0		5	0	0	0		0	0	0	0		5	0	0	0
	#21	Ø	A	B	C	#22	Ø	A	B	C	#23	Ø	A	B	C	#24	Ø	A	B	C
Alng l		6	0	0	5		6	0	0	5		6	0	0	5		5	0	0	4
Bet v l		6	5	0	6		6	3	0	5		6	5	0	5		6	0	0	4
Cor a l		4	0	0	0		4	0	0	0		0	0	0	0		0	0	0	0
Cor a 'll'		5	0	0	0		4	0	0	0		5	0	0	0		0	0	0	0
	#25	Ø	A	B	C	#26	Ø	A	B	C	#27	Ø	A	B	C	#28	Ø	A	B	C
Alng l		4	0	0	4		5	0	0	0		6	0	0	4		6	0	0	5
Bet v l		4	4	0	5		7	5	0	6		6	0	0	5		6	0	0	5
Cor a l		0	0	0	0		5	0	0	0		5	0	0	0		5	0	0	0
Cor a 'll'		0	0	0	0		5	0	0	0		5	0	0	0		5	0	0	0
	#29	Ø	A	B	C	#30	Ø	A	B	C	#31	Ø	A	B	C	#32	Ø	A	B	C
Alng l		6	0	0	5		7	0	0	5		6	0	0	5		5	0	0	4
Bet v l		6	5	0	6		7	0	0	6		7	0	0	6		7	5	0	7
Cor a l		5	0	0	0		5	0	0	0		4	0	0	0		0	0	0	0
Cor a 'll'		5	0	0	0		6	0	0	0		5	0	0	0		0	0	0	0
	#33	Ø	A	B	C	#34	Ø	A	B	C	#35	Ø	A	B	C	#36	Ø	A	B	C
Alng l		4	0	0	4		5	0	0	4		5	0	0	4		6	0	0	5
Bet v l		7	5	0	7		6	4	0	5		7	5	0	6		7	5	0	6
Cor a l		3	0	0	0		0	0	0	0		4	0	0	0		4	0	0	0
Cor a 'll'		3	0	0	0		0	0	0	0		4	0	0	0		5	0	0	0
	#37	Ø	A	B	C	#38	Ø	A	B	C										
Alng l		6	0	0	5		5	0	0	5										
Bet v l		7	4	0	6		5	0	0	5										
Cor a l		6	0	0	0		0	0	0	0										
Cor a 'll'		6	0	0	0		0	0	0	0										

Table 3. The reactivity pattern of 38 patient sera deduced from the solid phase absorption experiments

Specificity	Number of sera	Percentage of all sera
B, AB, ABC1, ABC2, ABC1C2	16	42.1
AB, ABC1, ABC2, ABC1C2	6	15.8
AB	5	13.2
B, AB	4	10.2
B, ABC1, ABC2, ABC1C2	2	5.3
A, B, AB, ABC1, ABC2, ABC1C2	2	5.3
AB, ABC1	1	2.6
B, BC1, BC2, AB, BC1C2, ABC1, ABC2, ABC1C2	1	2.6
B, AB, ABC2	1	2.6

All the serum samples were analysed in a likewise manner and Table 3 summarizes the specificity patterns deduced and the number of samples which exhibit a given pattern. The pattern observed for sample number 8 was by far the most frequent and only two sera may contain IgE which apart from other specificities exhibit distinct A reactivity. However, this assignment is very uncertain due to the weakness of the radiostaining of *Aln g I* after absorption with Bv and Ca. The specificity patterns ABC1C2, ABC1 and ABC2 cannot be verified in the same way as the other patterns and their existence primarily are deduced from intensity ranking and therefore the patterns are less certain than any of the other patterns observed.

Discussion

The majority of the proteins extractable from pollens of alder, birch and hazel seem to be present as at least immunochemically homologous variants in all the extracts [19,20,29]. The major allergens *Aln g I*, *Bet v I* and *Cor a I* exhibit a degree of similarity with respect to: molecular weight, pI variation, amino acid composition, NH₂-terminal or complete amino acid sequence, hydrophathy profile and secondary structure propensity [13–29] which strongly indicate that the fundamental three-dimensional structure of these molecules are the same [49], and the differences between the allergens presumably originate from individual but small perturbations of the general structure.

Rabbit antibodies raised towards isolated tree pollen major allergens exhibit a strong crossreactivity towards the allergens of the related species [19,29]. Although the patients IgE response originate from a fundamentally different immunization scenario, there is no reason to

assume that the IgE antibodies should exhibit any restriction with respect to their epitope specificity. Therefore it is likely that the IgE antibodies will be able to react with many, if not all, of the available epitopes on the major allergens and due to the similarity between the allergens, it is probable that the IgE antibodies may crossreact with two or more of the major allergens. Recently Lind *et al.* [9,10] demonstrated that the specificity profile of patients IgE response towards the major allergens of house dust mites, *Der p I*, *Der f I* and *Der m I* exhibit both species specific and common reactivity. Further, Borch *et al.* [15] showed that the IgE response towards *Co a I* and 'II' seem to be crossreactive.

The specificity pattern of the patients IgE, deduced from the solid phase absorption experiments, strongly indicate that the patients' cross sensitivity towards pollens from trees of the Fagales order originate from IgE which react with epitopes common to all the major allergens and/or combinations of reactivity towards epitopes common between at least two major allergens (Table 3). None of the 38 sera exhibited an absorption pattern compatible with an exclusive species specific response. The majority of the sera (68.4%) seemed to contain IgE antibodies which react towards both apparent *Bet v I* specific epitopes and combinations of apparent common epitopes (Table 3). Only two sera, apart from common reactivities, seem to exhibit specific reactivity towards an allergen (*Aln g I*) different from *Bet v I* (Table 3). Further, all the common specificity patterns observed included a reactivity towards *Bet v I*, i.e. IgE reacting with *Bet v I* and at least one of the other major allergens, indicating that birch pollen are the primary, if not the only, sensitization source for these patients. Clinical data, i.e. hyposensitization, of these patients with extracts of either birch pollen or combinations of alder, birch and hazel pollen extracts seem to substantiate the above conclusion, since treatment with birch pollen extract alone provide the same clinical efficacy as did the mixture of the extracts [2–4].

Acknowledgment

The skilful technical assistance of Annette Giselsson is gratefully acknowledged.

References

- 1 Dirksen A, Østerballe O. Common components in pollen extracts. *Allergy* 1980; 35:611–16.
- 2 Petersen BN, Janniche H, Munch EP, Wihl J-Å, Ipsen H, Løwenstein H. Immunotherapy with partially purified and standardized tree pollen extracts. I. Clinical results from a three-year double blind study of patients treated with pollen

- extracts either of birch or combinations of alder, birch and hazel. *Allergy* 1988; 43:353-62.
- 3 Wihl J-Å, Ipsen H, Petersen BN, Munch EP, Løwenstein H. Immunotherapy with partially purified and standardized tree pollen extracts. II. Results of skinprick tests and nasal provocation tests from a three-year double blind study of patients treated with pollen extracts of birch or combinations of alder, birch and hazel. *Allergy* 1988; 43:363-9.
 - 4 Ipsen H, Schwartz B, Wihl J-Å, Petersen BN, Munch EP, Janniche H, Løwenstein H. Immunotherapy with partially purified and standardized tree pollen extracts. III. Specific IgE response to the major allergens of alder, birch and hazel pollen during immunotherapy. *Allergy* 1988; 43:370-7.
 - 5 Möller C, Dreborg S. Cross-reactivity between deciduous trees during immunotherapy. I. In vivo results. *Clin Allergy* 1986; 16:135-44.
 - 6 Möller C, Dreborg S, Einarsson R. Immunotherapy to deciduous tree pollens: specific IgE and IgG antibody patterns. *Clin Allergy* 1987; 17:551-562.
 - 7 Matthiesen F, Løwenstein H. Group V allergens in grass pollens. II Investigation of group V allergens in pollens from 10-grasses. *Clin Exp Allergy* 1991; 21:309-20.
 - 8 Leiferman KM, Gleich GJ. The cross-reactivity of IgE antibodies with pollen allergens. I. Analysis of various species of grass pollens. *J Allergy Clin Immunol* 1976; 58:129-39.
 - 9 Lind P, Ingeman L, Brouvez M. Demonstration of species-specific sensitization to major allergens of *Dermatophagoides* species by solid phase absorption of human IgE antibodies. *Scand J Immunol* 1987; 25:1-10.
 - 10 Lind P, Hansen OC, Horn N. The binding of mouse hybridoma and human IgE antibodies to the major faecal allergen, *Der p 1*, of *D. pteronyssinus*. Relative binding site location and species specificity studied by solid-phase inhibition assays with radiolabelled antigen. *J Immunol* 1988; 140:4256-62.
 - 11 Chapman MD, Sutherland WM, Platts-Mills TAE. Recognition of two *Dermatophagoides pteronyssinus* specific epitopes on antigen P1 by using monoclonal antibodies: Binding to each epitope can be inhibited by serum from house dust mite-allergic patients. *J Immunol* 1984; 133:2488-95.
 - 12 Heyman PW, Chapman MD, Platts-Mills TAE. The antigen *Der f 1* from the dust mite *Dermatophagoides farinae*: structural comparison with *Der p 1* from *Dermatophagoides pteronyssinus* and epitope specificity of murine IgG and human IgE antibodies. *J Immunol* 1986; 137:2841-7.
 - 13 Belin L. Separation and characterization of birch pollen antigens with special reference to the allergenic components. *Int Arch Allergy* 1972; 42:329-42.
 - 14 Viander M, Fräki P, Djupsund BM, Laine, S. Antigens and allergens in birch pollen extracts. *Allergy* 1979; 34:289-302.
 - 15 Borch MS, Skålhegg BA. Some characteristics of allergen components in an aqueous extract from hazel pollen (*C. avellana*). *Allergy* 1980; 35:194-5.
 - 16 Apold J, Florvaag E, Elsayed S. Comparative studies on tree pollen allergens. I. Isolation and partial characterization of a major allergen from birch pollen (*Betula verrucosa*). *Int Arch Allergy Appl Immunol* 1981; 64:439-47.
 - 17 Florvaag E, Elsayed S, Apold J. Comparative studies on tree pollen allergens. II. Isolation of alder (*Alnus incana*) pollen allergens. Purification and some characteristics of the major allergen pI 4.78. *Int Arch Allergy Appl Immunol* 1982; 67:49-56.
 - 18 Ipsen H, Løwenstein H. Isolation and immunochemical characterization of the major allergen of birch pollen (*Betula verrucosa*). *J Allergy Clin Immunol* 1983; 72:150-9.
 - 19 Ipsen H, Bøwadt H, Janniche H, Petersen BN, Munch EP, Wihl J-Å, Løwenstein H. Immunochemical characterization of reference alder (*Alnus glutinosa*) and hazel (*Corylus avellana*) pollen extracts and the partial immunochemical identity between the major allergens of alder, birch and hazel pollens. *Allergy* 1985; 40:510-18.
 - 20 Ipsen H, Larsen JN. Detection of antigen specific IgE antibodies in sera from allergic patients by SDS-PAGE immunoblotting and crossed radioimmuno-electrophoresis. In: Bjerrum O, Heegaard NHH, eds. Handbook of immunoblotting of proteins. Boca Raton: CRC Press, 1985; vol 2:159-66.
 - 21 Florvaag E, Elsayed S, Hammer ASE. Comparative studies on tree pollen allergens. XIII. Partial characterization of the alder (*Alnus incana*) pollen extract by two dimensional IEF/SDS-PAGE electrophoresis combined with electrophoretic transfer and immunoautoradiography. *Int Arch Allergy Appl Immunol* 1986; 80:26-32.
 - 22 Vik H, Elsayed S. Comparative studies on tree pollen allergens. X. Further purification and N-terminal amino acid sequence analysis of the major allergen of birch pollen (*Betula verrucosa*). *Int Arch Allergy Appl Immunol* 1986; 80:17-25.
 - 23 Vorch MS, Sletten K, Hagen AM. Chemical and immunochemical characterization of a major allergen (*Cor a 1*) from hazel pollen (*Corylus avellana*). XVI Nordic Congress on Allergology 1987; Abstr VI-19, p. 65.
 - 24 Florvaag E, Holen E, Vik H, Elsayed S. Comparative studies on tree pollen allergens. XIV. Partial characterization of the birch (*Betula verrucosa*) and hazel (*Corylus avellana*) pollen extracts by IEF and 2-D SDS-PAGE combined with electrophoretic transfer and IgE immunoautoradiography. *Ann Allergy* 1988; 61:393-400.
 - 25 Breiteneder H, Pettenburger K, Bito A, Valenta R, Kraft D, Rumpold H, Scheiner O, Breitenbach M. The gene coding for the major birch pollen allergen *Bet v 1*, is highly homologous to a pea disease resistance response gene. *EMBO J* 1989; 8:1935-8.
 - 26 Jarolim E, Tjekl M, Rohac M, Scherka G, Breitenbach M, Scheiner O, Kraft D, Rumpold H. Monoclonal antibodies against birch pollen allergens. Characterization by immunoblotting and the use for single step affinity purification of the major allergen *Bet v 1*. *Int Arch Allergy Appl Immunol* 1989; 90:54-60.
 - 27 Wheeler AW, Deards M, Fox B, Hickman BE, Spackman VM. Induction of polyclonal and monoclonal antibodies with a single allergen separated from birch pollen extract by

- SDS-PAGE and nitrocellulose blots. *Int Arch Allergy Appl Immunol* 1989; 90:378-81.
- 28 Marc-Series I, Boutin Y, Vrancken ER, Hébert J. Mapping of *Bet v* I epitopes by using murine monoclonal antibodies. *Int Arch Allergy Appl Immunol* 1990; 92:226-32.
 - 29 Ipsen H, Hansen OC. The NH₂-terminal amino acid sequences of the immunochemically partial identical major allergens of alder (*Alnus glutinosa*) *Aln g* I, birch (*Betula verrucosa*) *Bet v* I, hornbeam (*Carpinus betulus*) *Car b* I and oak (*Quercus alba*) *Que a* I pollens. *Mol Immunol* 1991; 28:1279-88.
 - 30 Löwenstein H. Immunological partial identity and in vitro inhibitory effect of two major pollen allergens to whole pollen extract of four grasses. *Int Arch Allergy Appl Immunol* 1978; 57:379-83.
 - 31 Chakraborty S, Löwenstein H, Ekramoddoullah AKM, Kisil FT, Schon AH. Detection of cross-reactive allergens in Kentucky blue grass pollen and six other grasses by crossed radioimmuno-electrophoresis. *Int Arch Allergy Appl Immunol* 1985; 78:300-4.
 - 32 Esch RE, Klapper DG. Cross-reactive and unique grass group I antigenic determinants defined by monoclonal antibodies. *J Allergy Clin Immunol* 1987; 79:489-95.
 - 33 Esch RE, Klapper DG. Isolation and characterization of a major cross-reactive grass group I allergenic determinant. *Mol Immunol* 1989; 26:557-61.
 - 34 Matthiesen F, Schumacher M, Löwenstein H. Characterization of the major allergen of *Cynodon dactylon* (Bermuda grass) pollen. *J Allergy Clin Immunol* 1991; 88:763-4.
 - 35 Cottam GP, Moran DM, Slandring R. Physicochemical and immunochemical characterization of allergenic proteins from rye grass (*Lolium perenne*) pollen prepared by a rapid and efficient purification. *Meth Biochem J* 1986; 234:305-10.
 - 36 Perez M, Ishioka GY, Walker LE, Chesnut RW. cDNA cloning and immunological characterization of the Rye grass allergen *Lol p* I. *J Biol Chem* 1990; 265:16210-15.
 - 37 Griffith IJ, Smith PM, Pollack J, Theerakulpisut P, Avjioglu A, Davies S, Hough T, Singh MB, Simpson RJ, Ward LD, Knox RB. Cloning and sequencing of *Lol p* I, the major allergenic protein of rye grass pollen. *FEBS Lett* 1991; 279:in press.
 - 38 Singh MD, Hough T, Theerakulpisut P, Avjioglu A, Davies S, Smith PM, Taylor P, Simpson RJ, Ward LDF, McCluskey J, Pay R, Knox RB. Isolation of cDNA encoding a newly identified major allergenic protein of rye grass pollen: Intracellular targeting to the amyloplast. *Proc Natl Acad Sci USA* 1991; 88:1384-8.
 - 39 Haavik S, Paulsen BS, Wold JK. Glycoprotein allergens in pollen of timothy. II. Isolation and characterization of a basic glycoprotein allergen. *Int Arch Allergy Appl Immunol* 1985; 78:260-8.
 - 40 Mohapatra SS, Hill R, Astwood J, Ekramoddoullah AKM, Olsen E, Silvanovich A, Hattom T, Kisil FT, Schon AH. Isolation and characterization of a cDNA clone encoding an IgE-binding protein from Kentucky blue grass (*Poa pratensis*) pollen. *Int Arch Allergy Appl Immunol* 1990; 91:362-8.
 - 41 Silvanovich A, Astwood J, Zhang L, Olsen E, Kisil F, Schon AH, Mohapatra S, Hill R. Nucleotide sequence analysis of three cDNAs coding for *Poa p* IX isoallergens of Kentucky blue grass pollen. *J Biol Chem* 1991; 266:1204-10.
 - 42 Thomas WR, Stewart GA, Simpson RJ, Chua KY, Plozza TM, Dilworth RJ, Nisbeth N, Turner KJ. Cloning and expression of DNA coding for the major house dust mite allergen *Der p* I in *Escherichia coli*. *Int Arch Allergy Appl Immunol* 1988; 85:127-9.
 - 43 Chua KY, Stewart GA, Thomas WK, Simpson RJ, Dilworth RJ, Plozza TM, Turner KJ. Sequence analysis of cDNA coding for a major house dust mite allergen *Der p* I. Homology with cysteine proteases. *J Exp Med* 1988; 167:175-82.
 - 44 Chua KY, Doyle CR, Simpson RJ, Turner KJ, Stewart GA, Thomas WR. Isolation of cDNA coding for the major mite allergen *Der p* II by IgE plaque immunoassay. *Int Arch Allergy Appl Immunol* 1990; 91:118-23.
 - 45 Chua KY, Dilworth RJ, Thomas WR. Expression of *Dermatophagoides pteronyssinus* allergen, *Der p* II, in *Escherichia coli* and the binding studies with human IgE. *Int Arch Allergy Appl Immunol* 1990; 91:124-9.
 - 46 Høiby N, Axelsen NH. Crossed immunoelectrophoresis as modified for quantitative purposes. *Scand J Immunol* 1983; 17 Suppl 10:125-9.
 - 47 Wecke B, Søndergaard I, Lind P, Aukrust L, Löwenstein H. Cross radio immunoelectrophoresis (CRIE) for identification of allergens and detection of the antigenic specificity of patients' IgE. *Scand J Immunol* 1983; 17 Suppl 10:265-72.
 - 48 Østerballe O, Ipsen H, Wecke B, Löwenstein H. Specific IgE response towards allergenic molecules during perennial hyposensitization: a three year prospective double blind study. *J Allergy Clin Immunol* 1983; 71:40-6.
 - 49 Chothia C, Finkelstein AV. The classification and origin of protein folding patterns. *Ann Rev Biochem* 1990; 59:1007-39.